



A new estimate of carbon for Bangladesh forest ecosystems with their spatial distribution and REDD+ implications

Mukul, Sharif A.; Biswas, Shekhar R.; Rashid, A. Z. M. Manzoor; Miah, Md. Danesh; Kabir, Md. Enamul; Uddin, Mohammad Belal; Alamgir, Mohammed; Khan, Niaz Ahmed; Sohel, Md. Shawkat Islam; Chowdhury, Mohammad Shaheed Hossain ; Rana, Md. Parvez; Rahman, Syed Ajijur; Khan, Mohammed Abu Sayed Arfin; Hoque, Muhammad Al-Amin

Published in:

International Journal of Research on Land-use Sustainability

Publication date:

2014

Document version

Publisher's PDF, also known as Version of record

Citation for published version (APA):

Mukul, S. A., Biswas, S. R., Rashid, A. Z. M. M., Miah, M. D., Kabir, M. E., Uddin, M. B., Alamgir, M., Khan, N. A., Sohel, M. S. I., Chowdhury, M. S. H., Rana, M. P., Rahman, S. A., Khan, M. A. S. A., & Hoque, M. A-A. (2014). A new estimate of carbon for Bangladesh forest ecosystems with their spatial distribution and REDD+ implications. *International Journal of Research on Land-use Sustainability*, 1, 33-41.

A new estimate of carbon for Bangladesh forest ecosystems with their spatial distribution and REDD+ implications

Sharif A. Mukul^{1,2,3,*}, Shekhar R. Biswas⁴, A. Z. M. Manzoor Rashid⁵, Md. Danesh Miah⁶, Md. Enamul Kabir⁷, Mohammad Belal Uddin⁵, Mohammed Alamgir^{6,8}, Niaz Ahmed Khan⁹, Md. Shawkat Islam Sohel¹, Mohammad Shaheed Hossain Chowdhury¹⁰, Md. Parvez Rana¹¹, Syed Ajijur Rahman^{12,13}, Mohammed Abu Sayed Arfin Khan^{5,14}, Muhammad Al-Amin Hoque^{2,15}

¹School of Agriculture and Food Sciences, The University of Queensland, Brisbane, QLD 4072, Australia

²School of Geography, Planning and Environmental Management, University of Queensland, Brisbane, QLD 4072, Australia

³Centre for Research on Land-use Sustainability, Maijdi, Noakhali 3800, Bangladesh

⁴Department of Biology, York University, 955 Oliver Road, Thunder Bay, ON, Canada P7B5E1

⁵Department of Forestry and Environmental Science, School of Agriculture and Mineral Sciences, Shahjalal University of Science and Technology, Sylhet 3114, Bangladesh

⁶Institute of Forestry and Environmental Sciences, University of Chittagong, Chittagong 4331, Bangladesh

⁷Forestry and Wood Technology Discipline, Life Science School, Khunla University, Khulna 9208, Bangladesh

⁸Centre for Tropical Environmental and Sustainability Sciences, School of Earth and Environmental Sciences, James Cook University, Cairns, QLD 4870, Australia

⁹Department of Development Studies, University of Dhaka, Dhaka 1000, Bangladesh

¹⁰Department of Forest Science, Faculty of Agriculture, Shinshu University, 8304 Minamiminowa-Mura, Nagano-Ken, Japan

¹¹School of Forest Sciences, University of Eastern Finland, P.O. Box 111, FI – 80101 Joensuu, Finland

¹²Center for International Forestry Research (CIFOR), Bogor, Barat 16680, Indonesia

¹³Forest and Landscape, University of Copenhagen, 1958 Frederiksberg, Denmark

¹⁴Department of Disturbance Ecology, University of Bayreuth, 95440 Bayreuth, Germany

¹⁵Department of Geography and Environment, Jagannath University, Dhaka 1100, Bangladesh

*Corresponding author: sharif_a_mukul@yahoo.com / s.mukul@uq.edu.au

Received: June 1, 2014; Accepted: July 15, 2014

Communicated by: Dr. Tapan Sarker, Griffith University, Australia

ABSTRACT

In tropical developing countries, reducing emissions from deforestation and forest degradation (REDD+) is becoming an important mechanism for conserving forests and protecting biodiversity. A key prerequisite for any successful REDD+ project, however, is obtaining baseline estimates of carbon in forest ecosystems. Using available published data, we provide here a new and more reliable estimate of carbon in Bangladesh forest ecosystems, along with their geo-spatial distribution. Our study reveals great variability in carbon density in different forests and higher carbon stock in the mangrove ecosystems, followed by in hill forests and in inland Sal (*Shorea robusta*) forests in the country. Due to its coverage, degraded nature, and diverse stakeholder engagement, the hill forests of Bangladesh can be used to obtain maximum REDD+ benefits. Further research on carbon and biodiversity in under-represented forest ecosystems using a commonly accepted protocol is essential for the establishment of successful REDD+ projects and for the protection of the country's degraded forests and for addressing declining levels of biodiversity.

Key words: carbon budget, forest conservation, forest ecosystem, REDD+, Bangladesh.

Introduction

Globally, about 60% of the carbon is stored in

Highlights

- We estimate 251.8 million Mg of carbon stored in Bangladesh forest ecosystems, with nearly 49.4% stored in the mangrove forests alone;
- We find 179.1 million Mg Carbon in forest biomass and 72.7 million Mg carbon in soil;
- The hill forests have the highest potential for forest carbon enhancement and REDD+ in the country.

forests, with about 12-20% of anthropogenic greenhouse gas (GHG) emissions being attributable to forest degradation and loss (Baccini et al. 2012; Houghton et al., 2012; Paoli et al. 2010; Clark et al. 2001). In the tropical region, reducing emissions from deforestation and forest degradation (REDD+) is a new climate change mitigation mechanism that aims to promote carbon sequestration by compensating tropical countries for conserving their forests (Angelsen et al. 2009; Parker et al. 2009). REDD+, in

recent years, has been increasingly recognized for its potential to jointly address the issue of climate change mitigation and forest conservation in the tropics (Phelps et al. 2012). Since 2009, REDD+ programs have generated estimated carbon investments of about US\$2 billion worldwide (Budiharta et al. 2014; Peters-Stanley et al. 2013). The successful implementation of REDD+ projects, however, relies largely on the robust estimation of carbon stored in forest ecosystems, with demonstrated benefits to local biodiversity and people (Saatchi et al. 2011; Paoli et al. 2010).

REDD+ programs require participating countries to quantify GHG emissions and removals from forests in a manner that is reliable, transparent, and as accurate as possible (Phelps et al. 2012; Gibbs et al. 2007). In many countries, however, large uncertainties still remain surrounding the emission estimates that arise from inadequate data on carbon density in local forests and unreliable estimates of national deforestation rates (Baccini et al. 2012). Unfavorable policy environments and technical challenges of measuring, reporting, and verifying GHG emissions from forests also remain critical in many developing countries (Harris et al. 2012; Baker et al. 2010). Developing countries that wish to participate in REDD+ programs need to construct a reference emissions level, which can be used as a benchmark for estimating emission reductions to be achieved by local REDD+ projects (Harris et al. 2012). National-level distribution of carbon in aboveground forest biomass and soil based on improved data sets can be very useful and act as initial steps for successful REDD+ projects (Alamgir and Turton 2014; Mukul 2014; Shin et al. 2008). A national forest monitoring system and standardized protocols for measuring GHG emissions from forests are also important for the implementation of REDD+ projects in tropical developing countries (Blom et al. 2010; Angelsen et al. 2009).

Here, we provide a new estimate of carbon in Bangladesh forest ecosystems based on published literature. We compile and review related studies

that report carbon stocks in biomass and soil in Bangladesh forest ecosystems. In doing so, we first provide relevant information on land use and forests in Bangladesh within the broader context of climate change. We then categorize the forests into three major categories (i.e. evergreen to semi-evergreen hill forests, littoral mangrove forests, and deciduous Sal forests) based on vegetation and ecology, and estimated carbon in biomass and soil for each category, with relevant maps showing their geo-spatial distribution. We believe our study is an initial step to recognize the role played by Bangladesh forests in climate change mitigation. Our study uses scientifically rigorous, reliable, and locally available forest inventories and will be useful for the advancement of national REDD+ programs and forest-based climate change mitigation project developments.

Land-use, land use change and forestry (LULUCF) in Bangladesh

Agriculture is the main land-use in Bangladesh, comprising nearly 65% of the country's land area, followed by forests (17.5%), urban/built-up areas (7.9%), and water (Figure 1; Dey et al. 2012). Forests occupy about 2.53 million hectares in Bangladesh (Khan et al. 2007). Of these, 1.53 million hectares (10.53% of the country's total land) are under the jurisdiction of the Forest Department (FD) (Alamgir and Turton 2014; Table 1). The remaining areas are unclassified state forests (0.73 million hectares) under the control of district administration and homestead forests (0.27 million hectares) owned by smallholder landowners (Khan et al. 2007). Most of the forests in the country are regenerated secondary forests (53%), followed by primary (30%) and planted (17%) forests (Mongabay 2014).

Bangladesh lost about 2.8% (~58,000 ha) of its forest coverage between 2000 and 2012 (Hansen et al. 2014, 2013), and still has one of the highest rates of deforestation in Asia (Poffenberger 2000). Most of

Table 1. Forest lands under the Forest Department's control in Bangladesh

Forest type	Area (million hectare)	Percentage (%)	
		# country's forest area	# country's land area
Hill forests	0.67	44.1	4.5
Mangrove forests	0.60	39.6	4.1
Mangrove plantation	0.13	8.5	0.9
Sal forests	0.12	7.9	0.8
Total	1.52	100	10.3




Figure 1. (a) Major land-use/land cover in Bangladesh, (b) tree-based land-use/land cover, and (c) major forest types managed by the Forest Department.

the country's state-owned forests are degraded in nature (Biswas and Choudhury 2007). Rural people's dependence on forests and diverse stakeholder engagement make forest management in the country very complex and challenging (Rashid et al. 2013; Mukul et al. 2012).

The highly diverse hill forests are situated mainly in the Chittagong and Sylhet division and are evergreen to semi-evergreen in nature (Khan et al. 2007; Figure 2, a, b). The Sundarban mangrove forest is the world's largest contiguous mangrove forest and is situated in the Khulna district (Figure 2, c). It is dominated by Sundri (*Heritiera fomes*) and is renowned for being the home of the globally endangered Bengal tiger. The mangrove plantations (Figure 2, d) are located on the newly accreted islands of coastal Bangladesh and are dominated mainly by Keora (*Sonneratia apetala*). The deciduous Sal forests (Figure 2, e) are located on relatively plain lands in the central districts of Bangladesh and are dominated mainly by Sal (*Shorea robusta*). Although managed by smallholders and rural landowners, the homestead forests represent one the most productive systems in Bangladesh (Kabir and Webb 2008; Figure 2, f).

Forestry in the context of climate change mitigation in Bangladesh

About 20% of Bangladesh's GHG emissions are estimated to derive from LULUCF activities, although the contribution of the country to global GHG

emissions is rather low, being less than 0.2% (WRI 2014). Per capita CO₂ emissions are also one of the lowest in the world and were estimated at 0.37 metric tons in 2011 (World Bank 2014).

There is no complete inventory for forest carbon estimation in Bangladesh. The latest assessment by the Food and Agriculture Organization of the United Nations (FAO) (i.e. Global Forest Resources Assessment 2010) reports 81 million Mg (Megagram; 10³ kg) carbon in living forest biomass in the country (FAO 2010). By contrast, Alamgir and Turton (2014) reported carbon density between 49-121 Mg ha⁻¹ in the country's forests depending on the condition of the vegetation (e.g. open canopy versus closed canopy). A large variability exists also in the national-level estimates of forests' carbon density, which is mainly attributable to differences in methods and sampling strategies (see Table 2). Using carbon densities reported in the literature, the carbon stocks in FD-managed forests is estimated to be between 98.8 and 240.2 million Mg (Table 2). The estimated carbon storage in soil (up to 30cm depth) in the country's forests is about 92.9 million Mg (FAO 2007, 1977). The highest soil carbon density is in the mangrove forests due to the greatest concentration of organic carbon in soil and is the lowest in the mangrove plantations (see Table 3). The plainland Sal forests, due to their degraded nature, also have relatively lower levels of soil carbon density and stock.



Figure 2. Different forest types in Bangladesh, from top left: (a) hill forest in the Chittagong Hill Tracts, (b) hill forest in Sylhet, (c) mangrove forest in the Sundarbans, (d) a mangrove plantation in coastal Bangladesh, (e) Sal forest in central Bangladesh, and (f) homestead forest in southern Bangladesh (Photo credits: Sharif A. Mukul).

Table 2. National-level estimates of biomass carbon in FD-managed forests in Bangladesh

Source	Carbon density (Mg ha ⁻¹)*	Carbon stock (Million Mg)	Remark
Saatchi et al. 2011	70.5	107.2	Based on satellite data
Gibbs and Brown 2007	158	240.2	Based on forest inventory
Gibbs et al. 2007	65	98.8	Based on harvest data
IPCC 2006	93	141.4	Based on harvest data
DeFries et al. 2002	137	208. 2	Based on harvest data
Brown 1997	92	139.8	Based on forest inventory
Mean	102.6	155.9	

*using median value when providing a range.

Table 3. Soil carbon storage in different forest types in Bangladesh

Forest type	Carbon density* (Mg ha ⁻¹)	Carbon stock (Million Mg)	Remark
Hill forests	49.5	33.2	SOC– 1.5%, and bulk density - 1.1
Mangrove forests	88.2	52.9	SOC – 2.1%, and bulk density - 1.4
Mangrove plantation	19.6	2.6	SOC – 0.45%, and bulk density - 1.5
Sal forests	34.5	4.2	SOC – 1.0%, and bulk density - 1.15
Total		92.9	-

Sources: FAO (2007, 1977)

Estimate of carbon in Bangladesh forest ecosystems

We estimate 179.1 million Mg carbon in biomass and 72.7 million Mg carbon in soil (up to 30cm depth) in Bangladesh forest ecosystems that are currently under the FD's jurisdiction. These figures, however, may overestimate the actual carbon storage in Bangladesh forest ecosystems, recognizing the fact that many of the country's forests are degraded in nature. Tables 4 and 5 below show the reported carbon density in different forest types in

Bangladesh that are used for these estimates. Here, we only consider the studies that are from primary and old growth secondary forests and do not include studies from monoculture, agroforestry, and/or plantation forests. The number of studies on soil organic carbon (n=11) is relatively higher than the studies on biomass carbon (n=7). There are only a few studies on plainland Sal forests and there are no studies on mangrove plantations. The majority of the studies have been conducted in the hill forests of the Chittagong and Chittagong Hill Tracts.

Table 4. Carbon in biomass in different forest types of Bangladesh

Forest type	Carbon density* (Mg ha ⁻¹)	Source
Hill forests	103.4	Mukul 2014
	115.6	Ullah and Al-Amin 2012
	73.6	Alamgir and Al-Amin 2007
	92.0	Shin et al. 2007
Mangrove forests [‡]	98.9	Rahman et al. 2014
	126.7	Donato et al. 2011
Sal forests	153.9	Kibria and Saha 2011

*using median value when providing a range.

[‡]includes mangrove plantations.**Table 5.** Soil organic carbon in different forest types of Bangladesh

Forest type	Carbon density* (Mg ha ⁻¹)	Source
Hill forests	51.0	Mukul 2014
	23.1	Barua and Haque 2013
	50.5	Ullah and Al-Amin 2012
	54.0	Miah et al. 2009
	80.1	Chowdhury et al. 2007
	97.7	Shin et al. 2007
	39.7	Islam et al. 2001
Mangrove forests [‡]	33.6	Rahman et al. 2014
	43.9	Donato et al. 2011
Sal forests	58.5	Kibria and Saha 2011
	38.1	Isalm and Weil 2000

*using median value when providing a range.

[‡]includes mangrove plantations.

The average ecosystem carbon density in Bangladesh forests is estimated at about 175.5 Mg ha⁻¹ (Table 6). Despite its degraded nature, we find the highest carbon density in the Sal forests (202.2 Mg ha⁻¹), which could be due to a limited number of studies

(n=1) on biomass carbon and bias in sampling. This could be the same case for soil carbon in mangrove forests. Mangrove forests constitute the highest share (54%) of the country's biomass carbon stock, followed by hill forests (36%) and the Sal forests

(Figure 3). Soil organic carbon stock is, however, highest in the hill forests (38.6 million Mg), comprising nearly 53% of the country’s total carbon stock in soil (Figure 4). Based on the studies, we find

higher uncertainty in soil carbon density in the hill forests (47.1%) followed by the Sal forests (29.9%) and mangrove forests (Figure 5).

Table 6. Carbon density in major forest types in Bangladesh

Forest type	Carbon density (Mg ha ⁻¹)		
	Biomass	Soil	Total
Hill forests	96.1 (±17.86)	57.6 (±27.13)	153.7
Mangrove forests [‡]	112.8 (±19.66)	38.8 (±7.28)	151.5
Sal forests	153.9 (0)	48.3 (±14.42)	202.2
Mean	120.9	48.2	169.1

[‡]includes mangrove plantation.




Figure 3. Carbon stock in biomass in major forest types of Bangladesh (a), and their relative contribution (b).




Figure 4. Soil organic carbon storage in major forest types of Bangladesh (a), and their relative contribution (b).




Figure 5. Map showing the density and spatial distribution of carbon in Bangladesh forest ecosystems; (a) total ecosystem carbon, (b) biomass carbon and uncertainty, and (c) soil carbon and uncertainty.

Concluding remarks

Altogether, we find 251.8 million Mg carbon stored in Bangladesh's forest ecosystems, with nearly 49.4% stored in the mangrove forests alone. Our estimate, however, is based on a limited number of available field inventories conducted into Bangladesh forest ecosystems, limiting the reliability of our estimates. Further studies should cover underrepresented forest ecosystems in the country using a commonly accepted protocol and sampling strategy. The mainland Sal forests represent the most degraded ecosystem in the country, while the mangrove forests are relatively intact due to limited human influence (Donato et al. 2011; Rahman et al. 2010). The hill forests located in the Chittagong, Chittagong Hill Tracts, and Sylhet are severely degraded due to illegal logging and shifting cultivation (Mukul et al. 2014; Khan et al. 2007). Local dependence on forests and unsustainable resource collection are also common in the hill forests (Miah et al. 2014).

The spatial distribution of forests is not uniform in the country, scattered mostly in the few districts of southern, central, and coastal Bangladesh (see Figure 5). The quality and quantity of forests has also decreased over the past decades (Biswas and Chowdhury 2007). Although not properly recognized, the potential found in Bangladesh's forests in terms of climate change mitigation through sequestering atmospheric carbon is still undeniable (Rahman and Akter 2013; Brown et al. 1993). To fully utilize this opportunity, the carbon sequestration potential of the country's forests should be integrated with REDD+ and other available carbon payment schemes (Sohel et al. 2009).

Due to diverse stakeholder engagement, coverage and the current state of forests, the hill forests of the country have the highest potential for forest carbon enhancement and REDD+. Tackling deforestation is the cheapest option for reducing human-induced GHG emissions and thereby for addressing climate change (Houghton 2013). Restoration of degraded forests using locally available species is also an ideal strategy for forest carbon enhancement in the country (Miah et al. 2009). Protection of degraded forests to allow natural regeneration, establishment of plantation in barren lands, and avoiding deforestation are some other key strategies that could also aid forest carbon enhancement (Niels et al. 2002). Creation of protected areas in forests with fair and equitable access for local people and their involvement in forest management and governance are also critical (Chowdhury et al. 2014; Rashid et al. 2013). Understanding rural livelihoods and economic cost-benefits are core to successful forest management in the country (Biswas et

al. 2009). To be socially acceptable, the value of unit carbon for REDD+ projects, therefore, should be sufficient to compensate the foregone monetary value associated with changes to local and indigenous people's traditional forest use in Bangladesh.

References

- Alamgir, M., Al-Amin, M., 2007. Organic carbon storage in trees within different Geopositions of Chittagong (South) forest division, Bangladesh. *Journal of Forestry Research*, 18: 174–180.
- Alamgir, M., Turton S.M., 2014. Climate change and organic carbon storage in Bangladesh forests. In: Tuteja, N., Gill, S.S. (eds.). *Climate change and plant abiotic stress tolerance*. Wiley-VCH Verlag.
- Angelsen, A., Brockhus, M., Kanninen, M., mSills, E., Sunderlin, W.D., Wertz-Kanounnikoff, S. (eds). 2009. *Realising REDD+: National strategy and policy options*. Centre for International Forestry Research, Bogor, Indonesia.
- Baccini, A., Goetz, S.J., Walker, W.S., Laporte, N.T., Sun, M., Sulla-Menashe, D., Hackler, J., Beck, P.S.A., Dubayah, R., Friedl, M.A., Samanta, S., Houghton, R.A., 2012. Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps. *Nature Climate Change*, 2: 182-185.
- Baker, D.J., Richards, G., Grainger, A., Gonzalez, P., Brown, S., DeFries, R., Held, A., Kelndorfer, J., Ndunda, P., Ojima, D., Skrovseth, P., Souza Jr, C., Stolle, F., 2010. Achieving forest carbon information with higher certainty: A five-part plan. *Environmental Science & Policy*, 13: 249-260.
- Barua. S.K., Haque, S.M.S., 2013. Soil characteristics and carbon sequestration potentials of vegetation in degraded hills of Chittagong, Bangladesh. *Land Degradation & Development*, 24: 63-71.
- Biswas, S.R., Choudhury, J.K., 2007. Forests and forest management practices in Bangladesh: the question of sustainability. *International Forestry Review*, 9: 627-640.
- Biswas, S.R., Mallik, A.U., Choudhury, J.K., Nishat, A., 2009. A unified framework for the restoration of Southeast Asian mangroves - bridging ecology, society and economics. *Wetlands Ecology and Management*, 17: 365-383.
- Blom, B., Sunderland, T., Murdiyarso, D., 2010. Getting REDD to work locally: lessons learned from integrated conservation and development projects. *Environmental Science & Policy*, 13: 164-172.
- Brown, S., 1997. *Estimating biomass and biomass change of tropical forests: a primer* FAO Forestry Paper no. 134, Rome, Italy.
- Brown, S., Iverson, L.R., Prasad, A., Liu, D., 1993. *Geographical distributions of carbon in biomass*

- and soils of tropical Asian forests. *Geocarto International*, 4: 45-59.
- Budiharta, S., Meijaard, E., Erskine, P.D., Rondinini, C., Pacifici, M., Wilson, K.A., 2014. Restoring degraded tropical forests for carbon and biodiversity. *Environmental Research Letters*, 9: 114020.
- Chowdhury, M.S.H., Biswas, S., Halim, M. A., Haque, S.M.S., Muhammed, N., Koike, M., 2007. Comparative analysis of some selected macronutrients of soil in orangeorchard and degraded forests in Chittagong Hill Tracts, Bangladesh. *Journal of Forestry Research*, 18: 27-30.
- Chowdhury, M.S.H., Gudmundsson, C., Izumiyama, S., Koike, M., Nazia, N., Rana, M.P., Mukul, S.A., Muhammad, N., Redowan, M., 2014. Community attitudes towards forest conservation programs through collaborative protected area management in Bangladesh. *Environment, Development and Sustainability*, 16: 1235-1252.
- Clark, D.A., Brown, S., Kicklighter, D.W., Chambers, J.Q., Thomlinson, J.R., Ni, J., Holland, E.A., 2001. Net primary production in tropical forests: an evaluation and synthesis of existing field data. *Ecological Applications*, 11: 371-384.
- DeFries, R.S., Houghton, R.A., Hansen, M.C., Field, C.B., Skole, D., Townshend, J., 2002. Carbon emissions from tropical deforestation and regrowth based on satellite observations for the 1980s and 1990s. *Proc. Natl. Acad. Sci., USA*, 99: 14256-14261.
- Dey, N.C., Saha, R., Islam, M.A., Iqbal, M.A., 2012. Assessing land use changes and land degradation in Bangladesh. *Stamford Journal of Environmet and Human Habitat*, 1: 60-70.
- Donato, D.C., Kauffman, J.B., Murdiyarso, D., Kurnianto, S., Stidham, M., Kanninen, M., 2011. Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience*, 4: 293-297.
- FAO (Food and Agriculture Organization of the United Nations), 1977. *Soil map of the world, Volume VII: South Asia*. FAO, Rome, Italy.
- FAO, 2007. Digital soil map of the world. Available online at: <http://www.fao.org/geonetwork/srv/en/metadata.show?id=14116> (Date of access: April, 2014).
- FAO, 2010. *Global Forest Resources Assessment 2010*. FAO Forestry Paper 163, FAO, Rome, Italy.
- Gibbs, H.K., Brown, S., 2007. Geographical distribution of biomass carbon in tropical southeast Asian forests: an updated database for 2000. Oak Ridge National Laboratory, Oak Ridge, TN, USA. Available online at: <http://cdiac.ornl.gov/epubs/ndp/ndp068/ndp068b.html> (Date of access: May, 2014).
- Gibbs, H.K., Brown, S., Niles, J.O., Foley, J.A., 2007. Monitoring and estimating tropical forest carbon stocks: making REDD a reality. *Environmental Research Letters*, 2: 045023.
- Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau, D., Stehman, S.V., Goetz, S. J., Loveland, T.R., Kommareddy, A., Egorov, A., Chini, L., Justice, C.O., Townshend, J.R.G., 2013. High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science*, 342: 850-853.
- Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau, D., Stehman, S.V., Goetz, S.J., Loveland, T.R., Kommareddy, A., Egorov, A., Chini, L., Justice, C.O., Townshend, J.R.G., 2013. Tree Cover and Tree cover Loss and Gain, Country Profiles. University of Maryland, Google, USGS, and NASA. Available online at: <http://www.globalforestwatch.org/country/BGD> (Date of access: April, 2014).
- Harris, N.L., Brown, S., Hagen, S.C., Saatchi, S.S., Petrova, S., Salas, W., Hansen, M.C., Potapov, P. V., Lutsch, A., 2012. Baseline map of carbon emissions from deforestation in tropical regions. *Science*, 336: 1573-1576.
- Houghton, R.A., 2013. The emissions of carbon from deforestation and degradation in the tropics: past trends and future potential. *Carbon Management*, 4: 539-546.
- Houghton, R.A., House, J.I., Pongratz, J., van der Werf, G.R., DeFries, R.S., Hansen, M.C., Quere, C.L., Ramankutty, N., 2012. Carbon emissions from land use and land-cover change. *Biogeosciences*, 9: 5125-5142.
- IPCC (Intergovernmental Panel on Climate Change), 2006. *IPCC Guidelines for National Greenhouse Gas Inventories*. Institute for Global Environmental Strategies, Japan.
- Islam, K.R., Ahmed, M.R., Bhuiyan, M.K., Badruddin, A., 2001. Deforestation effects on vegetative regeneration and soil quality in tropical semi-evergreen degraded and protected forests of Bangladesh. *Land Degradation & Development*, 12: 45-56.
- Islam, K.R., Weil, R.R., 2000. Land use effects on soil quality in a tropical forest ecosystem of Bangladesh. *Agriculture, Ecosystems and Environment*, 79: 9-16.
- Kabir, M.E., Webb, E.L., 2008. Can homegardens conserve biodiversity in Bangladesh? *Biotropica*, 40: 95-103.
- Khan, M.A.S.A., Uddin, M.B., Uddin, M.S., Chowdhury, M.S.H., Mukul, S.A., 2007. Distribution and status of forests in the tropic: Bangladesh perspective. *Proc. Pakistan Acad. Sci.*, 44: 145-153.
- Kibria, M.G., Saha, N. 2011. Analysis of existing agroforestry practices in Madhupur Sal forest: an assessment based on ecological and economic perspectives. *Journal of Forestry*

- Research, 22: 533-542.
- Miah, M. D., Uddin, M.F., Bhuiyan, M.K., Koike, M., Shin, M.Y., 2009. Carbon sequestration by the indigenous tree species in the reforestation program in Bangladesh - *Aphanamixis Polystachya* Wall. and Parker. *Forest Science and Technology*, 5: 62-65.
- Miah, M.D., Akther, S., Shin, M.Y., Koike, M., 2014. Scaling up REDD+ strategies in Bangladesh: a forest dependence study in the Chittagong Hill Tracts. *Forest Science and Technology*, 10: 148–156.
- Mongabay, 2014. Bangladesh: Environmental Profile. Available online at: <http://rainforests.mongabay.com/deforestation/archive/Bangladesh.htm> (Date of access: May, 2014).
- Mukul, S.A., 2014. Biodiversity conservation and ecosystem functions of indigenous agroforestry systems: Case study from three tribal communities in and around Lawachara National Park. Pp 171-179. In: Chowdhury, M.S.H. (ed.). *Forest Conservation in Protected Areas of Bangladesh: Policy and community development perspective*. Springer, Switzerland.
- Mukul, S.A., Herbohn, J., Rashid, A.Z.M.M., Uddin, M.B., 2014. Comparing the effectiveness of forest law enforcement and economic incentive to prevent illegal logging in Bangladesh. *International Forestry Review*, 16: 363-375.
- Mukul, S.A., Rashid, A.Z.M.M., Quazi, S.A., Uddin, M.B., Fox, J., 2012. Local peoples' response to co-management regime in protected areas: A case study from Satchari National Park, Bangladesh. *Forests, Tress and Livelihoods*, 21: 16-29.
- Niles, J.O., Brown, S., Pretty, J., Ball, A.S., Fay, J., 2002. Potential carbon mitigation and income in developing countries from changes in use and management of agricultural and forest lands. *Phil. Trans. R. Soc. Lond. A* . 360: 1621-1639.
- Paoli, G.D., Wells, P.L., Meijaard, E., Struebig, M.J., Marshall, A.J., Obidzinski, K., Tan, A., Rafiastanto, A., Yaap, B., Slik, J.W.F., Morel, A., Perumal, B., Wielaard, N., Husson, S., D'Arcy, L., 2010. Biodiversity conservation in the REDD. *Carbon Balance and Management* 5: 7.
- Parker, C., Mitchell, A., Trivedi, M., Mardas, N., 2009. *The Little REDD+ Book*. Global Canopy Programme, Oxford, UK.
- Peters-Stanley, M., Hamilton K., Yin D., 2012. Leveraging the landscape, state of the forest carbon markets 2012. *Ecosystem Marketplace*, Forest Trends, USA.
- Phelps, J., Webb, E.L., Adams, W.M., 2012. Biodiversity co-benefits of policies to reduce forest-carbon emissions. *Nature Climate Change*, 2: 497-503.
- Poffenberger, M., 2000. Communities and forest management in South Asia. IUCN, DFID, and Asia Forest Network, the Philippines.
- Rahman, M. M., Khan, M.N.I., Hoque, A.K.F., Ahmed, I., 2014. Carbon stock in the Sundarbans mangrove forest: spatial variations in vegetation types and salinity zones. *Wetlands Ecology and Management*, doi: 10.1007/s11273-014-9379-x.
- Rahman, M.M., Rahman, M.M., Guogang, Z., Islam, K.S., 2010. A review of the present threats to tropical moist deciduous Sal (*Shorea robusta*) forest ecosystem of central Bangladesh. *Tropical Conservation Science*, 3: 90-102.
- Rahman, M.S., Akter, S., 2013. Carbon forestry: scope and benefit in Bangladesh. *Journal of Forest Science*, 29: 249-256.
- Rashid, A.Z.M.M., Craig, D., Jeffery. M.I., Khan, N.A., 2013. Forest protected area governance in Bangladesh: a focus on the legal and policy framework. *Chinese Journal of Population Resources and Environment*, 11: 345-351.
- Rashid, A.Z.M.M., Craig, D., Mukul, S.A., Khan, N.A., 2013. A journey towards shared governance: status and prospects of collaborative management in the protected areas of Bangladesh. *Journal of Forestry Research*, 24: 599-605.
- Saatchi, S.S., Harris, N.L., Brown, S., Lefsky, M., Mitchard, E.T.A., Salas, W., Zutta, B.R., Buermann, W., Lewis, S.L., Hagen, S., Petrova, S., White, L., Silman, M., Morel, A., 2011. Benchmark map of forest carbon stocks in tropical regions across three continents. *Proc. Natl. Acad. Sci., USA*, 108: 9899-9904.
- Shin, M.Y., Miah, M.D., Lee, K.H., 2007. Potential contribution of the forestry sector in Bangladesh to carbon sequestration. *Journal of Environmental Management*, 82: 260-276.
- Shin, M.Y., Miah, M.D., Lee, K.H., 2008. Mitigation options for the Bangladesh forestry sector: implications of the CDM. *Climate Policy*, 8: 243-260.
- Sohel, M.S.I., Rana, M.P., Alam, M., Akhter, S., Alamgir, M., 2009. The carbon sequestration potential of forestry sector: Bangladesh context. *Journal of Forest Science*, 25: 157-165.
- Ullah, M.R., Al-Amin, M., 2012. Above and below-ground carbon stock estimation in a natural forest of Bangladesh. *Journal of Forest Science*, 58: 372-379.
- World Bank, 2014. World Development Indicators: Bangladesh. Available online at: <http://databank.worldbank.org/data/reports.aspx?source=2&country=BGD&series=&period=> (Date of access: May, 2014).
- WRI (World Resources Institute), 2014. Global Forest Watch: Bangladesh Climate Data. Available online at: <http://climate.globalforestwatch.org/countries/BGD> (Date of access: May, 2014).

